

# AmproMed™ P FOR POULTRY

## (amprolium)

### 9.6% Oral Solution

### Coccidiostat



Water-soluble treatment for coccidiosis  
Treats 800 Gallons at 0.012% level.

#### INDICATIONS

AmproMed P FOR POULTRY (amprolium) 9.6 % Oral Solution is intended for the treatment of coccidiosis in growing chickens, turkeys, and laying hens. If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

#### USE DIRECTIONS

Give amprolium at the 0.012% level (8 fl oz per 50 gallons) as soon as coccidiosis is diagnosed and continue for 3 to 5 days. (In severe outbreaks, give amprolium at the 0.024% level). Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time. Use as the sole source of amprolium.

To prepare 50 Gallons of Medicated Water

DOSAGE LEVEL	MIXING DIRECTIONS
0.024 %	Add 1 pint (16 fluid ounces) of AmproMed P FOR POULTRY (amprolium) 9.6% Oral Solution in about 5 gallons of water in a 50-gallon medication barrel. Stir, then add water to the 50-gallon mark. Stir thoroughly.
0.012 %	Follow same directions as above but use 1/2 pint (8 fluid ounces) of AmproMed P FOR POULTRY 9.6% Oral Solution.
0.006 %	Follow same directions as above but use 4 fluid ounces of AmproMed P FOR POULTRY 9.6% Oral Solution.

#### For Automatic Water Proportioners

For automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

DOSAGE LEVEL	AmproMed P FOR POULTRY 9.6 % ORAL SOLUTION PER GALLON OF STOCK SOLUTION
0.024 %	41 fl oz
0.012 %	20.5 fl oz
0.006 %	10.25 fl oz

Note: Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

#### WARNING

Keep this and all drugs out of reach of children. NOT FOR HUMAN USE.

#### PRECAUTIONS

**FOR ORAL USE IN ANIMLAS ONLY. MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention.

#### ADVERSE EVENTS

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse drug events, for technical assistance or to obtain a copy of the SDS contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at [www.fda.gov/reportanimalae](http://www.fda.gov/reportanimalae).

#### STORAGE

Store container in an upright position. Store between 5°C - 25°C (41°F - 77°F), with brief excursions to 40°C. Benzoic acid as Sodium Benzoate 0.1% added as preservative.

#### Restricted Drug (California) – Use Only as Directed

Approved by FDA under ANADA # 200-496

List Number	Pack Size	Case Size
1AMP024	1 Gallon	4

AmproMed™ is a trademark of Bimeda, Inc.  
Rev. 12/20 M-23-0366

# **AmproMed™ P FOR POULTRY**

**(amprolium)**  
**9.6% Oral Solution**  
**Coccidiostat**



**MANUFACTURED BY:**

Bimeda, Inc.  
Le Sueur, MN 56058  
[www.bimeda.com](http://www.bimeda.com)

**N.A. CORP. ADDRESS:**

Bimeda, Inc.  
One Tower Lane  
Oakbrook Terrace, IL 61081

For further information and queries, contact Bimeda at (888) 524-6332 or [US-info@Bimeda.com](mailto:US-info@Bimeda.com)

AmproMed™ is a trademark of Bimeda, Inc.  
Rev. 12/20 M-23-0366