

# MULTIDOSE

Fluke and Worm Drench

DATA  
SHEET



***The complete answer to all your dosing requirements, including Roundworms, Lungworms, Mature and immature flukes down to four weeks old.***

- ✓ 1.5% w/v Levamisole Hydrochloride which is active against stomach and Lung Worms.
- ✓ 2.25% w/v Rafoxanide which is active against Mature and Immature Flukes.
- ✓ The combination of Rafoxanide and Levamisole permits a three way activity against Flukes, Lung and Stomach worms in sheep and cattle.
- ✓ Multidose Fluke and Worm drench is active against *Fasciola Spp.*, *Nematodirus Spp.*, *Dictyocaulus Spp.* and *Ostertagia Spp.* In sheep and cattle.
- ✓ Multidose is also active against *Haemonchus Spp.* and nasal bots (*Oestrus Ovis*).

LIST No	UNIT PACKAGE	CASE SIZE
1MUL059	150ml	24
1MUL060	500ml	12
1MUL061	1L	12

See reverse for Administration & Dosage



# MULTIDOSE

Levamisole Hydrochloride and w/v Rafoxanide



## COMPOSITION

Each ml contains:

Levamisole HCl B.P	15mg (1.5%w/v)
Rafoxanide B.P (Vet)	22.5 mg (2.25%w/v)

## Reported Efficacies against Liver Fluke (*Fasciola hepatica*)

	4 week-old Immature Fluke	6 week-old Immature Fluke	Adult Liver Fluke
Rafoxanide*	45-98%	85-99%	99-100%

## DOSAGE & ADMINISTRATION

For oral use in cattle, sheep and goats only.

To be given orally at a dose rate of 7.5mg/kg bodyweight for Levamisole HCl & 11.25mg/kg body weight for Rafoxanide practical dosage recommendations are as follows:

### Cattle

Body weight (kg)	Dose (ml)	Dose per Pack
50	25	20
100	50	10
150	75	6.7
200	100	5
250	125	4
Over 300kg	150	3.3

### Sheep & Goats

Body weight (kg)	Dose (ml)	Dose per Pack
Up to 15kg	5	100
25	10	50
35	15	33.3
Over 35	20	25

## RECOMMENDED WITHOLDING PERIOD

Animals should not be slaughtered for human consumption during treatment or for 30 days thereafter.

Carefully estimate the live weight of the animal. Use properly calibrated dosing equipment.

Not to be given to lactating cows or ewes whose milk is used for human consumption

Not to be given to cows or ewes during lactation whose milk is used for human consumption.

## PRECAUTIONS:

- Store below 30°C
- For animal use only
- Shake well before use
- Keep out of reach of children

## MANUFACTURED UNDER LICENCE OF:

**Bimeda AMEA Ltd.**

1B The Hebert Building,  
The Park, Carrickmines,  
Dublin 18, Ireland.

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

[www.bimeda.co.ke](http://www.bimeda.co.ke)