

ADACYCLINE LA 20%

20% w/v injectable solution for veterinary use

COMPOSITION

Each ml contains Oxytetracycline 200mg

DESCRIPTION

Adacycline is a wide spectrum bacteriostatic antibiotic. It acts on gram-positive and gram-negative micro-organisms. It is particularly sensitive to *Streptococci*, *Clostridia*, *E.coli*, *Shigellae*, *Brucellae*, *Salmonellae*, *Leptospirae*, etc. In addition it is effective against some *Mycoplasmas*, *Rickettsiae*, *Chlamydiae*, some protozoa and large viruses. After administration of therapeutic dose of Adacycline, a high level of Oxytetracycline in blood is achieved within half an hour to 4 hours.

INDICATIONS

Treatment of diseases caused by Oxytetracycline-susceptible organism in cattle, sheep and goats. Diseases include Pneumonia and Shipping Fever Complex associated with *Pasteurella spp.* and *Haemophilus spp.*, Infectious Bovine Kerato-conjunctivitis (pinkeye) caused by *Moraxella Bovis*, Foot Rot and Diphtheria caused by *Fusobacterium Necrophorum*, bacterial Enteritis (scours) caused by *Escherichia Coli*, Wooden Tongue caused by *Actinobacillus Lignierisii*, *Leptospirosis* caused by *Leptospira Pomona*; wound infections and acute metritis caused by strains of staphylococci /streptococci organisms sensitive to Oxytetracycline.

For cattle; Bronchopneumonia and other infections of the respiratory and gastrointestinal tracts, *Metritis*, *Mastitis*, *Septicaemia*, puerperal infections and secondary bacterial infections etc.

For sheep and goats; Infections of respiratory, urogenital tract and hooves, mastitis, infected wounds, etc.

DOSAGE AND ADMINISTRATION

Administer Adacycline 20% L.A by deep intramuscular injection. The dose is 20 mg/kg body weight (1ml /10kg bodyweight). The product is recommended for a single dose administration. Maximum recommended dose at one site;

Cattle	20ml
Sheep and goats	5ml
Pigs	10ml

CONTRA- INDICATIONS

Adacycline inj. is not intended for cats, dogs and horses. It should not be given to animals in late pregnancy, animals with severe damages of liver and kidney and to animals over sensitive to Oxytetracycline.

SIDE-EFFECTS

Sometimes a temporary swelling on the injection site occurs.

WITHDRAWAL PERIOD

From last treatment, within 21 days for meat and edible tissues and within 7 days for milk.

WARNING

A maximum of 20ml may be injected at any one site for cattle and 5ml maximum for sheep and goats. If total volume exceeds the above mentioned quantity, it should be divided and injected to more sites. The drug must not be diluted.

STORAGE

Store in a cool dry place, away from light.

CATEGORY

P.O.M

PACKING

Bottles of 50 and 100ml

Manufactured by:

Chongqing Fantong Animal Pharm. Co. Ltd, Chongqing, China

For
BIMEDA LTD.



BIMEDA LTD.

Funzi Road, Industrial Area. P.O. Box 30620-00100,
Nairobi, Kenya
Tel: +254 20 6537622-6 Fax: +254 20 6537628