

DEWORMER


b Bimeda®

Global Excellence in Animal Health

VERMOFAS GOLD

FLUKE & WORM DRENCH

in cattle, sheep, and goats



Available packs:
125ml, 500ml, 1Litre

A bright yellow suspension containing 3% w/v Levamisole Hydrochloride B.P and Oxytocyanide 6% w/v B.P. (Vet)

Also used as ovicidal against fluke and roundworm eggs.

For **TOTAL**
control of
flukes and worms



Funzi Road, Industrial Area.
P.O. Box 30620 - 00100, Nairobi, Kenya.
Tel: 02060537622-6

www.bimeda.co.ke

VERMOFAS GOLD

FLUKE & WORM DRENCH

Indication:

For oral administration to cattle, sheep & goats at a dose rate of 7.5mg Levamisole Hydrochloride and 15mg Oxytoclozamide per kg bodyweight.

Contra-indications and Warnings.:

Milk for human consumption may only be taken from cows after 72 hours from the last treatment. Animals must not be slaughtered for human consumption during treatment.

Animals may be slaughtered for human consumption only after 28 days from the last treatment.

Levamisole activity is not affected by benzimidazole resistance.

**KEEP OUT OF REACH OF CHILDREN
PROTECT FROM LIGHT.
DO NOT STORE ABOVE 30° C
FOR ANIMAL TREATMENT ONLY.
SHAKE THE CONTAINER BEFORE
USE.**

Manufactured by:

Medisel (K) LTD,
General Kago Road,
P.O. BOX 540 - 01000.
Thika, Kenya.

Dosage instructions:

Cattle - 2.5ml per 10kg bodyweight

Bodyweight	Dose	Doses in this bottle
50kg	12.5 ml	80
100kg	25 ml	40
150kg	37.5 ml	26.5
200kg	50 ml	20
250kg	62.5 ml	16
300kg	75 ml	13

(Cattle over 300kg a further 2.5 ml per additional 10kg)

Sheep & goats – 0.5ml per 2kg bodyweight

Bodyweight	Dose	Doses in this bottle
10kg	2.5 ml	400
20kg	5 ml	200
30kg	7.5 ml	133
40kg	10 ml	100
50kg	12.5 ml	180
60kg	15 ml	66

(Sheep & goats over 60kg a further 0.5ml per additional 2 kg)

